

The Effect of Methanolic extract of *Eucalyptus sp.* Leaves in Growth of some Opportunistic Fungi

Qays Majeed Issa¹, Hussein Hamed², Mustafa Ahmed², Muntadar Adil², Lujain Bassam², Mouge Raad²

¹ Department of Pharmacognosy, College of Pharmacy, Al-Nahrain University, Baghdad, Iraq

² Students of Pharmacy College, Al-Nahrain University, Baghdad, Iraq

Corresponding Author Email: dr.qais.majeed@nahrainuniv.edu.iq

Abstract

The recent study aimed to examine the Methanolic extract of *Eucalyptus sp.* leaves against some opportunistic fungi. The leaves of *Eucalyptus sp.* were collected, dried and then grinded. Powder of leaves was extracted using methanol and then filtered using filter papers. The filtrate was then centrifuged and then the filtrate was concentrated using a rotary evaporator. A stock solution was used to determine the inhibitory activity of the extract against opportunistic fungi. The results of detecting the active substances in the crude

Introduction

The incidence of fungal infections has increased in the last ten years, with fungi infecting more than a billion people annually, of who about a million and a half are infected with opportunistic fungi that may sometimes cause death. Fungi usually infect patients with asthma, diabetes, and cancer [1], as well as patients undergoing organ transplantation and those who are treated with corticosteroids [2]. The developing resistance of microorganisms against appealing antimycotic medications is considered as one of the very important matters that preoccupy researchers and physicians. The treatment of pathogenic microorganisms which include, bacteria, fungi, viruses, and protozoa, is difficult because of the continuous development of resistance to these organisms [3]. In one of the recent statistics, researchers concluded that human deaths could reach ten million

annually by the year 2050 due to the resistance of pathogenic microorganisms to antibiotics, thus exceeding the number of people who die from cancer currently [4]. The antifungals discovered so far are usually partially effective in affecting the pathogenic fungi, and these antifungals also affect the host tissues [5]. The antifungals currently used are divided into four main categories: azoles, polyenes, flucytosines, and echinocandins [6].

There has become an urgent need to discover new medicines that are more effective and have fewer side effects. This has been done by exploiting natural plant products in discovering new medicines [6,7]. Bioactive constituents play an essential role in discovers of drugs by provide as constituents of importance for synthetic changes [8]. *Eucalyptus* is a genus of over 700 species of flowering trees, in the myrtle family (Myrtaceae) [9]. *Eucalyptus* is an affluent origin of phytochemical constituents as Tannins, Glycosides, Phenols, Flavonoids, Saponin, Alkaloids, kaempferol, Propanoids extracted from roots, steams, and leaves of the plant [10, 11]. So far, it has been revealed that there are more than 18 chemical compounds in the *Eucalyptus* plant. It was found that the compound Eucalyptol is present in 79.88% of the total chemical compounds found in this plant, which was found to be biologically active against various microorganisms [12].

Materials and Methods

Collection of Plant Material

Eucalyptus leaves were collected during November 2022 from the public gardens of Al-Kadhimiya city. The plant was diagnosed based on available plant diagnosis books. The plant leaves were washed well with tap water and then dried in the oven at a temperature of 50°C .Then the leaves were grinded using a blender to get a powder.

Extraction Method

The method of [13] was used to obtain the plant leaves extract. 200 grams of plant leaf powder put in a 1000 ml glass beaker, then mix up with 800 ml of concentrated methanol (99%), It was mixed well and then placed on a magnetic stirrer at a temperature (70°C) for (4) hours with slow motion of mixture using

magnetic stirrer. The mixture was first filtered using a piece of medical gauze, after which the filtrate was separated using a centrifuge (3000 rpm for 10 min.). The extract was put in rotary evaporator for (1) hour to remove the solvent, so as to obtain a stock solution which used as an antifungal substance .

Chemical Diagnosis of Some Active Ingredients in *Ecalyptus* leaves extract

Saponins detection

1ml of the crude extract of the plant was taken and placed in a graduated cylinder, 20 ml of distilled water was placed on it, then the cylinder was shaken well for 3 minutes, as this led to the emergence of foam with a height of 3 cm , which indicates that the crude extract contains Saponins in composition of the crude extract of the plant (14).

Glycosides Detection

One ml of crude extract of plant was added to 2 ml of Benedict's reagent in a test tube , the mixture was agitated well , the tube was placed in water bath at a temperature of 100°C for 5 minutes , after which the tube was cooled , a red precipitate appeared at the bottom of the tube , indicating the presence of glycosides in the composition of the crude extract (14).

Tannins Detection

1% of FeCl_3 was added to 0.5 ml of the crude extract of the plant in a test tube, a bluish-green precipitate appeared, indicating the presence of tannins in the composition of the crude extract (14).

Quinones Detection

Two ml of extract was taken, few drops of concentrated H_2SO_4 were added and appearance of red color indicates the presence of quinones (14).

Fungal isolates

Five genera of fungi were used in this study were *Aspergillus niger*, *Penicillium sp.*, *Rhizopus sp.*, *Alternaria sp.* and *Candida albicans* . They were isolated from plants and soils from Botanical garden of Pharmacognosy department - college of Pharmacy / Al-Nahrain University. All Fungi were isolated and purified using

culture medium of Potato Dextrose Agar (PDA). The phenotypic, microscopic and biochemical characteristics were relied upon to diagnose all isolated fungi (15).

Antifungal activity evaluation

The activity of the Methanolic extracts of plant leaves was examined by agar dilution method as declared by (16) to determination of minimum inhibitory concentrations (MICs) for fungi used in this study. A 6mm diameter culture disc from the edge of a newly growing fungal colony was transferred by using Cork Borer from each of the inspected isolates to the center of a plates containing fresh solid culture medium (SDA). After 7 days of incubation at a temperature of 28 °C, influence of various concentrations of the crude extract of plant leaves on the growth of the fungal cultures used in this study was examined by using a measuring ruler to determine the diameters of the fungal colonies for the purpose of calculating the percentage of inhibition using the following equation:

$$\text{Percentage of inhibition} = \frac{C - T}{C} \times 100$$

Where, C = colony diameter (mm) of the control

T = colony diameter (mm) of the test plate.

Results

Chemical diagnosis of some active ingredients in *Eucalyptus* leaves extract

Table (1): Chemical diagnosis of some active ingredients in *Eucalyptus* leaves extract

No.	Detection type	Result
1	Saponins detection	+
2	Glycosides detection	+
3	Tannins detection	+
4	Quinones detection	+

The results showed in table (2) that the Methanolic extract of *Eucalyptus sp.* Leaves had a clear effect on the growth of *A. niger*, starting from a concentration

of (0.5%), while it had the strongest effect at a concentration of (8%) may indicated the 100% complete inhibition of fungal growth.

Table (2): Inhibitory Activity of Methanolic extract of *Eucalyptus sp.* leaves on *Aspergillus niger* after incubation period 7 days, 28°C

Fungus	Extract concentration (%)	Colony diameter (cm)	Inhibition rate (%)
A. niger	control	7.5	0
	0.5	6	20
	1	3	60
	2	1	86.67
	4	0.6	92
	8	0	100

The Results showed in table (3) that the methanolic extract of *Eucalyptus sp.* leaves had a clear effect on the growth of the fungus *Fusarium sp.* starting from a concentration of (0.5%), while it had the strongest effect at a concentration of (8%) may led to 100% inhibition of fungal growth.

Table (3): Inhibitory Activity of methanolic extract of *Eucalyptus sp.* leaves on *Fusarium sp.* after incubation period 7 days, 30°C

Fungus	Extract concentration (%)	Colony diameter (cm)	Inhibition rate (%)
Fusarium sp.	control	6	0
	0.5	5.5	8.34
	1	4	33.34
	2	2	66.67
	4	0.8	86.67
	8	0	100

Results showed in table (4) that the crude methanolic extract of *Eucalyptus sp.* leaves had a clear effect on the growth of the fungus *Alternaria sp.* starting from a concentration of 0.5%, while it had the strongest effect at a concentration of 4% may indicate 100% inhibition of fungal growth by 100%.

Table (4): Inhibitory Activity of Methanolic extract of *Eucalyptus sp.* leaves on *Alternaria sp.* after incubation period 7 days, 30°C.

Fungus	Extract concentration (%)	Colon y diameter (cm)	inhibition rate (%)
Alternaria sp.	control	8	0
	0.5	6	25
	1	3	62.50
	2	1.5	81.25
	4	0	100
	8	0	100

Results in table (5) showed that the crude methanolic extract of *Eucalyptus sp.* leaves had a clear effect on the growth of the fungus *Rhizopus sp.* starting from a concentration of 0.5%, while it had the strongest effect at a concentration of 8% may led to the complete inhibition of fungal growth by 100%.

Table (5): Inhibitory Activity of Methanolic extract of *Eucalyptus sp.* leaves on *Rhizopus sp.* after incubation period 7 days, 30°C

Fungus	Extract concentration (%)	Colony diameter (cm)	Inhibition rate (%)
Rhizopus sp.	control	9	0
	0.5	8.5	5.56
	1	7	22.23
	2	3.5	61.12
	4	1	88.89
	8	0	100

Results showed in table (6) that the methanolic extract of *Eucalyptus sp.* leaves had a clear effect on the growth of the fungus *penicillium sp.* starting from a concentration of (0.5%) , while it had the strongest effect at a concentration of (8%) may led to the complete inhibition of fungal growth by (100%).

Table (6): Inhibitory Activity of Methanolic extract of *Eucalyptus sp.* leaves on *Penicillium sp.* after incubation period 7 days, 30°C

Fungus	Extract conc. (%)	Colony diameter (cm)	Inhibition rate (%)
Penicillium sp.	control	7	0
	0.5	6	14.29
	1	4	42.86
	2	2.5	64.28
	4	1	85.71
	8	0	100

Discussion

The fungal cell membrane is made up of ergosterol, which vary from the mammalian cell, which essentially contains cholesterol. Ergosterol is the main target when studying antimycotic drugs (17).

It has been found that flavonoids found in many plants are antimicrobial agents for a wide range of pathogenic microorganisms (18, 19) the mechanisms of action of antifungal drugs is to affect the functioning of the cytoplasmic membrane through preventing the biosynthesis of ergosterol in the fungal cell wall, this will lead to the leakage of cell components from the inside to the outside. Excessive production of reactive oxygen will also cause high oxidative stress to the fatty acids, amino acids, and nucleic acids present inside the fungal cell, and this will lead to an increase in the permeability of the cytoplasmic membrane (19).or may causing a malfunction in the functioning of mitochondria, antifungal medications cause a defect in the functioning of mitochondria . This defect may be due to a weakness in the respiratory process, or due to a weakness in the oxidation of fatty acids, or damage to the DNA of the mitochondria, or the reason may be a change in the permeability of the mitochondria. (19). Or through disrupting of DNA synthesis, cell division, protein synthesis, and cell wall formation. The antimycotic mechanism act based on distort of cell wall due to inhibition of the synthesis of β -glucans and chitin (19). Flavonoids affected the fungal filaments growth by reduce the G0/G1 phase and rising the G2/M phase. The inhibition of microtubule polymerization leads to inhibit the mitotic spindle formation and discouragement of cell division (20). The antimycotic agents enter inside the cell over active transport that arrives at inside the nucleus, hence inhibits protein synthesis, DNA, and RNA. Flavonols inhibit hyphal fungus *Cochliobolus lunatus* over the inhibition of DNA and RNA synthesis. Carvacrol, are disrupts the cellular plasma

membrane and impedes synthesis of the nucleic acids (19). Studies indicate that the volatile oil of the eucalyptus plant affects the physiological activity of fungi by inhibiting glucose breakdown, which in turn affects cell energy metabolism. (21). *E. camaldulensis* essential oil perfectly inhibited the mycelium of 5 strains of *Fusarium* spp. at a concentration scope (7 - 8 μ L/mL) after incubation for 5 days (22). In a study, results are revealed that phenol and terpene at (10%) concentration was notably best than others in growth inhibition of *F. oxysporum* which recorded 100 %, while alkaloid at (10 %) was notably less than others registered (63.66%) only (23). Another study showed that phenol extracted from *Eucalyptus* with a concentration of 10% inhibited the *F.oxysporum* fungus by 33.34%, as it was found that the antifungal properties depend on the lipophilic and hydrophilic properties.(24). These oils led to decay of fungal mycelium that revealed clear of cytoplasmic contents (25). Phenols effect on fungal cells due to these constituents sensitize the phospholipids of the fungal plasma membrane rising permeability of essential vital intracellular constituents (26).

Conclusion

The crude Methanolic extract of *Eucalyptus sp.* leaves was found to be a good source of antifungal agents. Significant antimycotic activity agent's up-against the investigated fungal isolates were as well noticed which is likely because of the existence of high bulk of saponins and phenolic compounds in the crude extract. This study highlights the possible utilize of bioactive constituents obtained from *Eucalyptus* extract as antimycotic agents in the industrial production of medicine in the future.

References

1. Bongomin F, Gago S, Oladele R, Denning, D. Global and Multi-National Prevalence of Fungal Diseases—Estimate Precision. *J. Fungi*. 2017; 3(4):57.
2. Lu M, Li T, Wan J, Li X., Yuan L., Sun S. Antifungal effects of phytocompounds on *Candida* species alone and in combination with fluconazole. *Int. J. Antimicrobial. Agents*.2017; 49:125–136.
3. Kannaiyan M, Meseret AG, Kanimozhi C, Thambidurai P, Ashokapuram Selvam S., Vinodhini R., *et al.* Prevalence of extended-spectrum beta-lactamase producing enterobacteriaceae members isolated from clinically suspected patients. *Asian J. Pharma. Clin. Res*.2018; 11:364.
4. Roemer T, Krysan DJ. Antifungal Drug Development: Challenges, Unmet Clinical Needs, and New Approaches. *Cold Spring Harbor Perspect. Med.* 2014; 4:019703.
5. Mickymaray, S, Al Aboody MS. *In vitro* antioxidant and bactericidal efficacy of 15 common spices: novel therapeutics for urinary tract infections? *Medicina*. 2019; 55:289.
6. Mickymaray, S, Alturaiki W. Antifungal Efficacy of Marine Macroalgae against Fungal Isolates from Bronchial Asthmatic Cases. *Molecules*. 2018; 23:3032
7. Moorthy K, Punitha T, Vinodhini R, Mickymaray S, Shonga A, Tomass Z *et al.* Efficacy of different solvent extracts of *Aristolochia krisagathra* and *Thottea ponmudiana* for potential antimicrobial activity. *J. Pharmacy Res*. 2015; 9:35–40.
8. Thawabteh A, Juma S, Bader M, Karaman D, Scrano L, Bufo SA *et al.* The Biological Activity of Natural Alkaloids against Herbivores, Cancerous Cells and Pathogens. *Toxins*. 2019; 11:656.
9. Macphail, M, Thornhill AH. "How old are the eucalypts? A review of the microfossil and phylogenetic evidence". *Australian Journal of Botany*. 2016;64 (8): 579.
10. Dixit A, Rohilla A, Singh V. *Eucalyptus globulus*: A new perspective in therapeutics. *Int J Pharm Chem Sci*. 2012; 1(4): 1678-83.

11. Boulekbache-Makhlouf L, Meudec E, Chibane M, *et al.* Analysis by high-performance liquid chromatography diode array detection mass spectrometry of phenolic compounds in fruit of *Eucalyptus globulus* cultivated in Algeria. *J Agric Food Chem* 2010; 58(24): 12615-24.
12. Emara S, Shalaby AE. Seasonal variation of fixed and volatile oil percentage of four *Eucalyptus* spp. related to lamina anatomy. *Afr J Plant Sci.* 2011; 5(6): 353-9.
13. Okogun JI. *Methods of Medicinal Plant Extract Preparation.* National Institute for Pharmaceutical Research and Development (NIPRD) 2000.
14. Kaur, Gupta S, Gautam P. Phytochemical analysis of *Eucalyptus* leaves extract. *J Pharmacogn Phytochem.* 2019; 8(1): 2442-2446
15. Forbes B, Sahm D, Weissfeld A. *Bailey and Scott,s : Diagnostic Microbiology.* 12th edition. St. Louis, Mo.: Elsevier Mosby, USA. 2007, pp.628-716.
16. Balouiri M, Sadiki M, Ibsouda S. Methods for *in vitro* evaluating antimicrobial activity: A review. *J Pharm Anal.*2016; 6(2):71-79.
17. Halat, D, Younes S, Mourad N, Rahal M. Allylamines, Benzylamines, and Fungal Cell Permeability: A Review of Mechanistic Effects and Usefulness against Fungal Pathogens. *Membranes (Basel).* 2022; 12(12): 1171.
18. Kumar S, Pandey A. Chemistry and Biological Activities of Flavonoids: An Overview. *Scientific World Journal.* 2013; 162750.
19. Al Aboody MS, Mickymaray S. Anti-Fungal Efficacy and Mechanisms of Flavonoids. *Antibiotics.*2020; 9(2): 45.
20. Hasim S, Coleman J. Targeting the fungal cell wall: current therapies and implications for development of alternative antifungal agents. *Future Med Chem.* 2019; 11(8): 869–883.
21. Zhou LJ, Li FR, Huang LJ, Yang ZR, Yuan S, Bai LH. Antifungal Activity of *Eucalyptus* Oil against Rice Blast Fungi and the Possible Mechanism of Gene Expression Pattern. *Molecules.* 2016; 21(5): 621.

22. Lopez-Meneses AK, Plascencia-Jatomea M, Lizardi-Mendoza J, Rosas-Burgos EC, Luque-Alcaraz AG, Cortez-Rocha MO. Antifungal and antimycotoxigenic activity of essential oils from *Eucalyptus globulus*, *Thymus capitatus* and *Schinus molle*. *Food Sci. Technol.* 2015; 35 (4).
23. Fradi, A and Al-Araji, A. Effect of *Eucalyptus camaldulensis* Terpens, Alkaloids and Phenols Against *Fusarium oxysporum*. *Iraqi Journal of Science* 2015; 56(4):2807-2810.
24. Knobloch K, Pauli A, Iberl N, Weigand, N and Weis, HM. Antibacterial and antifungal properties of essential oil components. *Journal of Essential Oil Research.* 1989; 1:119–128.
25. Hiroshi, N. Prevention of drug access to bacterial targets: permeability barriers and active efflux. *Science.* 1994; 264:382–388.
26. Simonetti G, Brasili E, Pasqua G. Antifungal Activity of Phenolic and Polyphenolic Compounds from Different Matrices of *Vitis vinifera* L. against Human Pathogens. *Molecules.* 2020; 25(16): 3748.